

Prof. Dott. CORRADINO CAMPISI
Professore Straordinario di Chirurgia Generale, angioproctologia
Direttore della UOS Dipartimentale di Chirurgia del Intestino - III
Specialista in Chirurgia Generale, vascolare, d'Urgenza e Pronto Soccorso
Università degli Studi di Genova

Abstract : Analisi dei risultati di un campione di studio relativo ai primi 50 casi trattati con “Biorisonanza” in pazienti affetti da Linfedema

C.Campisi, C.Eretta, D. Pertile, Da Rin Elisa, F.Boccardo - Genova

PER LA SOCIETÀ ITALIANA DI LINFANGIOLOGIA. S.I.L.

Lymphedema is a pathologic condition where the lymphatic flow is obstructed or proceeds very slowly: the result is a progressive swelling of the limbs.

A typical complication is recurrent acute lymphangitis (erysipelas) that is responsible for a worsening of the clinical picture (swelling and hardness of the limb).

This will let the lymphedema progress very fast, bringing the patient to invalidating conditions and (rarely) to tumoral degeneration (Stewart-Treves syndrome).

The continuous use of conservative treatment (bandaging, physiotherapy and compressive stockings) bring for sure an improvement of the lymphedema.

In selected cases, microsurgery of the lymphatics leads to an improvement if it is associated with conservative therapy, especially if performed in an early stage of the disease.

In our experience we also associated the Bioresonance together with the conservative and surgical therapy.

We chose a group of 50 patients affected from lymphedema (primary or secondary) of the arms or legs, previously treated with a combination of physiotherapy and surgery. We associated the use of the Bioresonance QRS Medithera and we observed an objective and subjective improvement of the clinical picture (volume and rate of recidives of erysipelas) in this group of patients.

Genova 22 Gennaio 2008

Prof. Dott. C. Campisi

